Fate Report for Case # P-18-0219

Fate

```
Summary Statement
              Fate P-18-0219
        Summary FATE:
       Statement: MW =
                              with < 500 \text{ and } < 1000
                          with MP < 25 °C
                   (E)
                   S = Reacts
                   Hydrolysis half-life = hr-da
                   VP < 1.0E-6 torr
                   at 25 °C (E)
                   BP > 400 \, ^{\circ}C \, (E)
                   H < 1.00E-8 (E)
                   POTW removal
                   (%) = PMN 90 via hydrolysis; then Hyd Pdt 90 via sorption
                   Time for
                   complete ultimate aerobic biodeg = Hyd Pdt > mo
                   Sorption to
                   soils/sediments = Hyd Pdt v.strong
                   PBT Potential: PMN P1B1; Hyd Pdt
                   *CEB FATE: Migration to ground water = Hyd Pdt negl
                   PMN
                   Material:
                   Overall wastewater treatment removal is 90% via hydrolysis
                   (hydrolysis half-life: hours to days).
                   PMN Material:
                   Low
                   Persistence (P1) is based on hydrolysis of the parent compound (hydrolysis
                   half-life: hours to days).
                   Low Bioaccumulation potential (B1) is
                   based on hydrolysis of the parent compound (hydrolysis half-life: hours to
                   Hydrolysis Product:
                   Overall wastewater treatment removal is
                   90% via sorption.
                   Sorption to sludge is strong based on data for
                                    polymers.
                   Air Stripping (Volatilization to air) is
                   negligible based on data for
                                                                    polymers.
                   Removal
                   by biodegradation in wastewater treatment is negligible based on data for
```

polymers. The aerobic aquatic biodegradation half-life is greater than months based on data for polymers. The anaerobic aquatic biodegradation half-life is greater than months based on the aerobic biodegradation half-life. The anaerobic biodegradation half-life is projected to be greater than or equal to the aerobic biodegradation half-life. Sorption to soil and sediment is very strong based on data for polymers. Migration to groundwater is negligible based on data for polymers. Hydrolysis Product: **High Persistence** (P3) is based on the aerobic and anaerobic biodegradation half-lives. Low Bioaccumulation potential (B1) is based on data for polymers and low water solubility, which inhibits bioavailability and biodegradation. Bioconcentration/Bioaccumulation factor to be put into E-Fast: N/A.



Physical Properties

Property	Measured/Calculated Value	EPI
Molecular Form:	1	
Molecular Wt.:		
% < 500:		
% < 1000:		

Property	Measured/Calculated	EPI
	Value	

Property	Measured Value	Method	Estimated Value	Method	EPI
Melting					
Point:					
Boiling			>400		
Point:					
BP					
Pressure:					
Vapor			< 0.000001		
Pressure:					
Water			Reacts		
Solubility:					
Log P:					
Log					
Kow:					
Log Koc:					
Log BCF:					
Henry's					
Law:					
nН•					

pH:	
pН	
Comment:	

Fate Analysis

Hydrolysis (t1/2,	Volatilization	Volatilization
da):	(t1/2)	(t1/2)
	- River (hr):	- Lake (da):
Atm Ox Potential	Atm Ox Potential	Atm Ox Potential
(t1/2)OH (hr):	(t1/2)O3	(t1/2) Total
	(hr):	(hr):
MITI Linear:	MITI	
	NonLinear:	
Biodeg Linear:	Biodeg	
	NonLinear:	
Biodeg Survey	Biodeg Survey	
ult:	Prim:	
STP (% removal)	STP (% removal)	
Total:	Biodeg:	

STP (% removal)	STP (% removal)	
Ads:	Air:	

Rationales

Removal in	
Wastewater	
Treatment:	
Atmospheric	
Oxidation:	
Hydrolysis:	
Photolysis:	
Aerobic	
Biodegradation:	
Anaerobic	
Biodegradation:	
Sorption	
to Soil and	
Sediment:	
Migration to	
Groundwater:	
Persistence - Air:	
Persistence	
- Water:	
Volatilization	
from Water:	
Soil:	
Sediment:	
Other:	
Standard:	
Bioaccumulation:	

PBT Ratings

2 1 1100011190			
Persistence	Bioaccumulation	Toxicity	PBT Comments
1	1		PMN
3	1		Hyd
			Pdt

Exposure-Based Testing

Exposure-Based	
Testing:	

Fate Ratings Removal in WWT/POTW (Overall):

Removal in 90;90 WWT/POTW PMN;Hyd Pdt (Overall):

Condition	Rating		Rating Do	escription		Comment
	Values	1	2	3	4	
WWT/POTW	;3	Low	Moderate	Strong	V. Strong	PMN;Hyd
Sorption:						Pdt
WWT/POTW	;4	Extensive	Moderate	Low	Negligible	, ,
Stripping:						Pdt
Biodegradation	;4	Unknown	High	Moderate	Negligible	, ,
Removal:		** •		-		Pdt
Biodegradation		Unknown	Complete	Partial		
Destruction:	4		XX / 1	3.6 (1		DANIII 1
Aerobic	;4	<= Dava	Weeks	Months	> Months	PMN;Hyd Pdt
Biodeg Ult: Aerobic		Days <=	Weeks	Months	NIOIIIIS	Put
Biodeg Prim:		_ Days	weeks	Months	Months	
Anaerobic	;4	<= Days	Weeks	Months	>	PMN;Hyd
Biodeg	,4	<- Days	WEEKS	Monuis	Months	Pdt
Ult:					Wioning	1 dt
Anaerobic		<=	Weeks	Months	>	
Biodeg Prim:		Days	,, 00115	1110114110	Months	
Hydrolysis (t1/2	2-3	<=	Hours	Days	>=	-N=C=O
at pH		Minutes		3	Months	
7,25C) A:						
Hydrolysis (t1/2		<= Minutes	Hours	Days	>=	
at pH 7,25C) B:					Months	
Sorption to	;1	V.	Strong	Moderate	Low	PMN;Hyd
Soils/Sediments:		Strong				Pdt
Migration to	;1	Negligible	Slow	Moderate	Rapid	PMN;Hyd
Ground						Pdt
Water:		NT 11 11 1	C1	N	D :1	
Photolysis A,		Negligible	Slow	Moderate	Rapid	
Direct: Photolysis B,		Magligible	Clarr	Moderate	Donid	
Indirect:		Negligible	Slow	Moderate	Rapid	
Atmospheric Ox		Negligible	Slow	Moderate	Rapid	
A, OH:		regugione	DIOW	Moderate	καρια	
11, 011.		Negligible	Slow	Moderate	Rapid	
		11051151010	510 11	1,10001010	Tapia	

Condition	Rating		Rating Description			
	Values	1	2	3	4	
Atmospheric Ox						
B, O3:						

Bio

Comments:

Bio The isocyanate groups are
Comments: expected to hydrolyze with a half-life of hours to days, to give the amine-terminated polymer.

Fate

Comments:

Fate Comments:

Comments/Telephone

Log

Artifact	Update/Upload	
	Time	